

Sea ice pCO₂ dynamics and related air-ice CO₂ fluxes during a flood-freeze cycle (Bellingshausen Sea, Antarctica).

N.-X. Geilfus (1), J.-L. Tison (2), G. Carnat (2), A.V. Borges (1), S. F. Ackley (3), and B. Delille (1)

(1) Chemical Oceanography Unit, Université de Liège, Belgium, (2) Glaciology Unit, Université Libre de Bruxelles, Belgium (3) Department of Earth and Environmental Science, University of Texas at San Antonio, U.S.A. (nxgeilfus@ulg.ac.be)

Sea ice covers about 7% of the Earth surface at its maximum seasonal extent. For decades sea ice was assumed to be an impermeable and inert barrier for air – sea exchange of CO₂ so that global climate models do not include CO₂ exchange between the oceans and the atmosphere in the polar region. However, uptake of atmospheric CO₂ over sea ice cover has been recently reported raising the need to further investigate pCO₂ dynamics in the marine cryosphere realm and related air-ice CO₂ fluxes.

We carried out direct measurements of pCO₂ within brines and related air-ice CO₂ fluxes (chamber method) of Antarctic first year pack ice during the “Sea Ice Mass Balance in Antarctica –SIMBA” drifting station experiment. This experiment was carried out on board the N.B. Palmer in September and October 2008 in the Bellingshausen Sea (Antarctica, 69 – 71°S; 90 – 95°E). In this area, significant snow loading promotes flooding of sea ice floes. In the course of the experiment we experienced a full cycle of cooling and warming of the air temperature with large changes in the snow cover thickness. Temperature and snow cover changes affected brine salinities, drastically increasing the instability of the brine column in the initial isothermal and porous sea ice. Cooling of the surface layer has significantly increased the surface layer brine salinity, triggering a downwards transfer of brines into the underlying porous ice. This downward transfer was likely counterbalanced by upwards (or lateral) transfer of sea water into the ice, a flooding-like process. While the sea-ice cover was undersaturated in CO₂ with respect to the atmosphere, convective processes significantly affected the partial pressure of CO₂ (pCO₂) of the brines, promoting the increase of pCO₂ and reducing the magnitude of related air-ice CO₂ transfer.